



Androgen dependent mechanisms of pro-angiogenic networks in placental and tumor development



Veronika M. Metzler^a, Simone de Brot^a, Robert S. Robinson^a, Jennie N. Jeyapalan^a, Emad Rakha^b, Thomas Walton^c, David S. Gardner^a, Emma F. Lund^a, Jonathan Whitchurch^d, Daisy Haigh^a, Jack M. Lochray^a, Brian D. Robinson^e, Cinzia Allegrucci^a, Rupert G. Fray^f, Jenny L. Persson^{g,h}, Niels Ødumⁱ, Regina R. Miftakhova^{h,j}, Albert A. Rizvanov^j, Ieuan A. Hughes^k, Rieko Tadokoro-Cuccaro^k, David M. Heery^d, Catrin S. Rutland^{a,**}, Nigel P. Mongan^{a,l,*}

^a Faculty of Medicine and Health Sciences, School of Veterinary Medicine and Health Sciences, University of Nottingham, LE12 5RD, UK

^b School of Medicine and Sciences, University of Nottingham, Nottingham City Hospital, NG5 1PB, UK

^c Department of Urology, Nottingham University Hospitals NHS Trust, NG5 1PB, UK

^d School of Pharmacy, University of Nottingham, NG7 2TQ, UK

^e Department of Pathology, Weill Cornell Medicine, New York 10065, USA

^f School of Biosciences, University of Nottingham, LE12 5RD, UK

^g Department of Translational Medicine, Lund University, Malmö, Sweden

^h Department of Molecular Biology, Umeå University, Sweden

ⁱ Department of Immunology and Microbiology, University of Copenhagen, Denmark

^j Kazan Federal University, Kazan, Republic of Tatarstan 420008, Russian Federation

^k Department of Paediatrics, University of Cambridge, Hills Rd, Cambridge CB2 0QQ, UK

^l Department of Pharmacology, Weill Cornell Medicine, New York 10065, USA

ARTICLE INFO

Article history:

Received 14 November 2016

Received in revised form

14 February 2017

Accepted 15 February 2017

Keywords:

Nuclear receptor

Epigenetics

Angiogenesis

VEGF

ABSTRACT

The placenta and tumors share important characteristics, including a requirement to establish effective angiogenesis. In the case of the placenta, optimal angiogenesis is required to sustain the blood flow required to maintain a successful pregnancy, whereas in tumors establishing new blood supplies is considered a key step in supporting metastases. Therefore the development of novel angiogenesis inhibitors has been an area of active research in oncology. A subset of the molecular processes regulating angiogenesis are well understood in the context of both early placentation and tumorigenesis. In this review we focus on the well-established role of androgen regulation of angiogenesis in cancer and relate these mechanisms to placental angiogenesis. The physiological actions of androgens are mediated by the androgen receptor (AR), a ligand dependent transcription factor. Androgens and the AR are essential for normal male embryonic development, puberty and lifelong health. Defects in androgen signalling are associated with a diverse range of clinical disorders in men and women including disorders of sex development (DSD), polycystic ovary syndrome in women and many cancers. We summarize the diverse molecular mechanisms of androgen regulation of angiogenesis and infer the potential significance of these pathways to normal and pathogenic placental function. Finally, we offer potential research applications of androgen-targeting molecules developed to treat cancer as investigative tools to help further delineate the role of androgen signalling in placental function and maternal and offspring health in animal models.

© 2017 Elsevier Ltd. All rights reserved.

* Corresponding author. Faculty of Medicine and Health Sciences, School of Veterinary Medicine and Health Sciences, University of Nottingham, LE12 5RD, UK.

** Corresponding author.

E-mail addresses: Catrin.rutland@nottingham.ac.uk (C.S. Rutland), Nigel.mongan@nottingham.ac.uk (N.P. Mongan).

1. Introduction

It has long been recognized that the placenta and tumors share important characteristics. These include mechanisms related to